

PanAfrican Medical Journal

Research

Sexual and reproductive health of adolescents in schools for people with disabilities

Open Access

CrossMark

Mercy Obasi¹, Stephen Manortey², Kofi Adesi Kyei^{3,4}, Michael Kwabeng Addo⁵, Sharon Talboys¹, Lynette Gay¹, Frank Baiden¹

¹Department of Public Health, University of Utah, West Africa Campus, Kpong, Ghana, ²Department of Radiography, School of Biomedical & Allied Health Sciences, Accra, Ghana, ³National Centre for Radiotherapy and Nuclear Medicine, Karle-Bu Teaching Hospital, Accra, Ghana

*Corresponding author: Kofi Adesi Kyei, Department of Radiography, School of Biomedical & Allied Health Sciences, Accra, Ghana

Key words: Adolescent, assessment, disability, sexual, reproductive health

Received: 26/02/2019 - Accepted: 24/03/2019 - Published: 14/06/2019

Abstract

Introduction: persons with disabilities have the same sexual and reproductive health (SRH) needs as the able-bodied people but they often face barriers to SRH information and services which are necessary for healthy and safe relationships, protection from HIV and other sexually transmitted infections (STIs). This study sought to assess the SRH services among adolescents with disabilities in four Special Needs Schools in Ghana. **Methods:** the study adopted a cross-sectional study design with a quantitative approach to data collection between the months of January to March, 2018. A structured and pretested questionnaire was used to collect data from adolescents with disabilities from selected schools in Ghana. Both descriptive and inferential statistics were performed using chi-square test and multivariate logistic regression. **Results:** majority of participants had hearing disability (52.1%). The average age at menarche among females was 13 years while the age at which puberty was attained among boys was 14 years. School teachers were the major source of information on SRH for the respondents (63.7%) followed by parents (12.2%). A majority (67.1%) of respondents had good knowledge of SRH. Factors which were significantly associated with knowledge level were age ($p=0.026$), religion ($p=0.024$), sources of information ($p<0.001$), guardians ($p=0.049$). **Conclusion:** the majority of participants had good knowledge of SRH, although their knowledge of contraceptive and access were poor. Only condoms were mostly known. There is the need for increased awareness on the availability of other contraceptives methods and the removal of barriers to contraceptive methods.

Pan African Medical Journal. 2019;33:299. doi:[10.11604/pamj.2019.33.299.18546](#)

This article is available online at: <http://www.panafrican-med-journal.com/content/article/33/299/full/>

© Mercy Obasi et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pan African Medical Journal – ISSN: 1937- 8688 (www.panafrican-med-journal.com)
Published in partnership with the African Field Epidemiology Network (AFENET). (www.afenet.net)

Updated 30 November 2023 12:51:39 by The AfriChild Centre Admin