

Venue-Based HIV  
Testing at Sex Work  
Hotspots to Reach  
Adolescent Girls and  
Young Women  
Living With HIV: A  
Cross-sectional  
Study in Mombasa,  
Kenya

We estimated the potential number of newly diagnosed HIV infections among adolescent girls and young women (AGYW) using a venue-based approach to HIV testing at sex work hotspots.

- [Venue-Based HIV Testing at Sex Work Hotspots to Reach Adolescent Girls and Young Women Living With HIV: A Cross-sectional Study in Mombasa, Kenya](#)

# Venue-Based HIV Testing at Sex Work Hotspots to Reach Adolescent Girls and Young Women Living With HIV: A Cross-sectional Study in Mombasa, Kenya

## IMPLEMENTATION SCIENCE

### OPEN

## Venue-Based HIV Testing at Sex Work Hotspots to Reach Adolescent Girls and Young Women Living With HIV: A Cross-sectional Study in Mombasa, Kenya

Huiting Ma, MSc;<sup>a</sup> Linwei Wang, MSc;<sup>a</sup> Peter Gichangi, MD, PhD;<sup>b,c</sup> Vernon Mochache, MD, MPH, PhD;<sup>d</sup> Griffins Manguro, MBChB;<sup>e</sup> Helgar K. Musyoki, MPH;<sup>f</sup> Parinita Bhattacharjee, MSc;<sup>g,h</sup> François Cholette, MPH;<sup>h,i</sup> Paul Sandstrom, PhD;<sup>h,i</sup> Marissa L. Becker, MD, MSc;<sup>e</sup> and Sharmistha Mishra, MD, MSc, PhD;<sup>h,i,k</sup> on behalf of the Transitions Study Team

**Background:** We estimated the potential number of newly diagnosed HIV infections among adolescent girls and young women (AGYW) using a venue-based approach to HIV testing at sex work hotspots.

Received for publication December 9, 2019; accepted March 15, 2020.  
From the <sup>a</sup>MAP-Centre for Urban Health Solutions, St. Michael's Hospital, Unity Health Toronto, Toronto, Canada; <sup>b</sup>Department of Human Anatomy, University of Nairobi, Nairobi, Kenya; <sup>c</sup>International Centre for Reproductive Health-Kenya, Mombasa, Kenya; <sup>d</sup>University of Maryland, Center for International Health, Education and Bioregistry, College Park, MD; <sup>e</sup>National AIDS & STI Control Programme, Nairobi, Kenya; <sup>f</sup>Key Populations Technical Support Unit, Partners for Health and Development in Africa, Nairobi, Kenya; <sup>g</sup>Centre for Global Public Health, University of Manitoba, Winnipeg, Canada; <sup>h</sup>National HIV and Reproductive Laboratory, <sup>i</sup>IC: With Infectious Diseases Research Centre, Public Health Agency of Canada, Winnipeg, Canada; <sup>j</sup>Department of Medical Microbiology and Infectious Diseases, University of Manitoba, Winnipeg, Canada; <sup>k</sup>Department of Medicine, University of Toronto, Toronto, Canada; <sup>l</sup>Institute of Medical Science, University of Toronto, Toronto, Canada; and <sup>m</sup>Institute of Health Policy, Management, and Evaluation, University of Toronto, Toronto, Canada.

Supported by the Canadian Institutes of Health Research operating Grant MOP13044 and foundation Grant FDN 13455. S.M. is supported by a Canadian Institutes of Health Research (CIHR) and Ontario HIV Treatment Network New Investigator Award. M.L.B. is supported by a CIHR New Investigator Award. The funders had no role in study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication. The authors have no funding or conflicts of interest to disclose.

H.M., L.W., and S.M. conceptualized and designed the study and developed the plan of analyses. M.L.B., S.M., H.K.M., and P.B. developed the study tools; M.L.B., P.G., G.M., H.K.M., and P.B. led the hotspot enumeration and Transitions study data collection. F.C. and P.S. led the serological testing and developed the reference testing algorithms. All authors contributed to interpretation of results and manuscript editing. S.M. and H.M. drafted the manuscript. H.M. conducted the analyses with input from L.W. and S.M.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site ([www.jaids.com](http://www.jaids.com)).  
Correspondence to: Sharmistha Mishra, MD, MSc, PhD, MAP-Centre for Urban Health Solutions, St. Michael's Hospital, University of Toronto, 200 Victoria Street, Toronto, ON M5B 1T8, Canada (e-mail: [sharmistha.mishra@utoronto.ca](mailto:sharmistha.mishra@utoronto.ca)).

Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

470 | [www.jaids.com](http://www.jaids.com)

J Acquir Immune Defic Syndr • Volume 84, Number 5, August 15, 2020

**Title:** Venue-Based HIV Testing at Sex Work Hotspots to Reach Adolescent Girls and Young Women Living With HIV: A Cross-sectional Study in Mombasa, Kenya

**Author (s):** Huiting Ma, Linwei Wang, Peter Gichangi, Vernon Mochache, Griffins Manguro, Helgar K. Musyoki, Parinita Bhattacharjee, François Cholette, Paul Sandstrom, Marissa L. Becker, and Sharmistha Mishra.

**Year of Publication:** 2020

**Access Resource Here:**

## INTRODUCTION

Adolescent girls and young women (AGYW) aged 15–24 years face a disproportionate risk of HIV acquisition in sub-Saharan Africa (SSA).<sup>1</sup> In Kenya, AGYW comprise 18.4% of the adult population but acquired 23.7% of new infections in 2017, such that, by 2018, an estimated 2.6% of AGYW in Kenya were living with HIV<sup>2–5</sup>; yet, most infections remain undiagnosed.<sup>4</sup> The most recent data available on AGYW suggest that, in 2012, only 25% of AGYW living with HIV were diagnosed and aware of their HIV status.<sup>4</sup> The consequence of undiagnosed HIV among AGYW